



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE SIMPLEST MODEL FOR ILLUSTRATING THE CONIC SECTIONS.

By LEONARD E. DICKSON, B. Sc., Fellow in Pure Mathematics, in the Chicago University.

The Greek geometers prior to Appollonius of Perga supposed that three cones were necessary in forming the conic sections. Thus, the ellipse was cut from an acute-angled cone; the parabola, from a right-angled cone; the hyperbola, from an obtuse-angled cone, each by a plane perpendicular to an edge. The very names, ellipse, parabola, and hyperbola, express the fact that the angle at the vertex of the cone is less than, equal to, or greater than a right angle; and thus that in the ellipse, the cutting plane *falls short* of the other nappe of the cone; in the parabola, is *parallel* to the edge of the cone; in the hyperbolia, *reaches over* to the other nappe of the cone.

Thus the earlier Greeks thought the ellipse peculiar to the acute-angled cone, the parabola to the right-angled cone, the hyperbola to the obtuse-angled cone.

But Appollonius of Perga showed in the year 250 B. C. that all three conics could be cut from single cone by varying the inclination of the cutting plane. Thus he cut a right-circular cone by a plane. Now if the angle between the cutting plane and the base of the cone be *less* than the angle made by an edge of the cone with the base, the section is an *ellipse*; if *equal*, a *parabola*; if *greater* an *hyperbola*.

The method of Appollonius, now used in our models consists in cutting a fixed cone by a revolving plane; my method consists in making the plane fixed and revolving the cone.

Nature furnishes us with a fixed horizontal plane, the surface of a liquid at rest. Filling partly full a hollow glass cone with some liquid (colored to make the effect distinct), we can make at will all the conic sections and their limiting positions. A definite amount of liquid will give a certain series of conics. By varying the quantity of liquid, we can form series of conics magnified in different degrees. Thus we can vary their curvature at will.

Suppose our cone is lying with an edge on a plane. If the plane be horizontal, the section is always a parabola, even if the cone be rolled about on the plane. If the plane is inclined to the horizon at an angle less than the angle at the vertex of the cone, then by rolling the cone about on the plane, the section passes successively through various forms of ellipses, then a parabola, then various forms of hyperbola, and again a parabola, and lastly the original ellipses.

This model has the great superiority over the ordinary wooden models, etc., in that we may readily study by it how the conics pass from one class into another through the limiting forms, the parabola, the circle, and the straight line.

Again, any school boy can make use of his conical ink-bottle, if no better model be at hand.

My method may be hapily extended to the construction of simple models for showing the various sections of any solid whatever.

[Read before the Texas Academy of Science in 1892.]